

UNCLASSIFIED

AD NUMBER
AD820848
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; JUL 1967. Other requests shall be referred to Commanding Officer, Fort Detrick, Attn: Technical Release Branch, Technical Information Division, Fort Detrick, MD
AUTHORITY
bdr1, d/a ltr, 29 Sep 1971

THIS PAGE IS UNCLASSIFIED

Best Available Copy

AD

AD820848

TECHNICAL MANUSCRIPT 402


PHOTOCONTROL OF ETHYLENE PRODUCTION
BY LETTUCE SEEDS AND BEAN HYPOCOTYLS

Frederick B. Abeles
Robert E. Holm
Harry E. Gahagan, III

JULY 1967

DEPARTMENT OF THE ARMY
Fort Detrick
Frederick, Maryland

Best Available Copy



**Best
Available
Copy**

Reproduction of this publication in whole or in part is prohibited except with permission of the Commanding Officer, Fort Detrick, ATTN: Technical Releases Branch, Technical Information Division, Fort Detrick, Frederick, Maryland, 21701. However, DDC is authorized to reproduce the publication for United States Government purposes.

DDC AVAILABILITY NOTICES

Qualified requesters may obtain copies of this publication from DDC.

Foreign announcement and dissemination of this publication by DDC is not authorized.

Release or announcement to the public is not authorized.

DISPOSITION INSTRUCTIONS

Destroy this publication when it is no longer needed. Do not return it to the originator.

The findings in this publication are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

DEPARTMENT OF THE ARMY
Fort Detrick
Frederick, Maryland 21701

TECHNICAL MANUSCRIPT 402

PHOTOCONTROL OF ETHYLENE PRODUCTION
BY LETTUCE SEEDS AND BEAN HYPOCOTYLS

Frederick B. Abeles

Robert E. Holm

Harry E. Gahagan, III

STATEMENT #2 UNCLASSIFIED

This document is subject to special export controls and each transmittal to foreign governments or foreign nationals may be made only with prior approval of *Co. Fort Detrick*

*attn: Technical Release Council
Technical Information Division
Fort Detrick, Md. 21701*

Crops Division
BIOLOGICAL SCIENCES LABORATORY

Project 1L013001A91A

July 1967

ACKNOWLEDGMENT

We gratefully acknowledge the expert technical assistance of Mr. Leonard E. Forrence during these investigations.

ABSTRACT

Red light of about 660 m μ has been shown to promote the germination of Lactuca sativa var. Grand Rapids seeds and to cause the unfolding of the hypocotyl hook of Phaseolus vulgaris L. var. Black Valentine. In both cases, the effect of red light was reversed by far-red light at about 730 m μ .

Ethylene promoted the germination of lettuce seeds and caused a negative curvature of bean hooks. Red light stimulated ethylene production from lettuce seeds but it inhibited ethylene production from the bean hooks. Lettuce seed germination and bean hook unfolding seem to be examples in which phytochrome promotes or inhibits the formation of ethylene that, in turn, appears to control the physiological response.

I. INTRODUCTION

Physiological processes in plants influenced by ethylene include abscission, dormancy, epinasty, flowering, fruit ripening, growth inhibition, membrane permeability, root initiation, and swelling. Burg¹ has reviewed these and other effects of the gas. Occasionally these processes are also influenced by other growth regulators, and evidence is rapidly accumulating that some of the effects of indoleacetic acid,²⁻⁸ herbicides,⁹⁻¹² gibberellin,³ and abscisic acid³ are mediated through ethylene.

This paper presents evidence that ethylene production is under photo-control and that some of the effects of light on plants may be mediated through ethylene.

II. MATERIALS AND METHODS

Seeds of Lactuca sativa var. Grand Rapids were germinated in gas-collection bottles described previously⁴ containing 10 ml of 1.5% agar. Each bottle contained 110 mg of seeds (about 100). During the 16-hour imbibition period, the seeds were exposed to far-red light.

Seedlings of Phaseolus vulgaris var. Black Valentine were grown in the dark in vermiculite for 6 to 7 days. Methods for harvesting, handling, and determining the hook opening followed those described by Klein et al.¹³ Ten hypocotyl hooks were set, base down, in 10 ml of 1.5% agar in 125-ml Erlenmeyer flasks sealed with rubber vaccine caps.

Far-red light was provided by a 150-watt incandescent bulb and was filtered by a Corning 165-mm square filter (Glass No. 2600 CS 7-69). For red light, the filter was a far-red Corning filter (Glass No. 3961 CS 1-56). Except for exposure to red or far-red light, all manipulations were performed under a green light (Corning filter CS 5-75) that had no effect on seed germination or hook opening. A temperature of 24 ± 2 C was maintained during the experiments.

Levels of ethylene surrounding the experimental material were determined by gas chromatography.⁴

III. RESULTS

The effect of light on hook opening is shown in Table 1. Hooks were excised from 6- to 7-day-old bean seedlings, placed on moist paper toweling to permit the escape of ethylene from wounded sections, then placed base down in agar and given light treatments (Table 1). The position of the bottles was shifted hourly during the light exposure so that incident energy was distributed as uniformly as possible. As observed earlier by others,¹³ the hook opening was stimulated by red light and inhibited by far-red light. Red light inhibited ethylene production from the hypocotyl sections and far-red light stimulated ethylene production. Data in the column headed Δ Curvature are the final curvature minus the initial curvature as determined by the methods of Klein et al.¹³ A positive value indicates that the hook opened and a negative value indicates that the hook closed. Addition of 1 ppm ethylene to the gas phase surrounding the hooks resulted in a Δ curvature of -40 degrees even in the presence of red light.

TABLE 1. EFFECT OF LIGHT ON HOOK OPENING OF
BLACK VALENTINE BEANS

Treatment		Δ Curvature, a/ degrees	C_2H_4 in Gas Phase, ppm	
0 to 5 hours	5 to 10 hours		0 to 12 hours	12 to 18 hours
dark	dark	-11	0.48	0.12
red	dark	+20	0.36	0.11
red	red	+46	0.38	0.07
red	far-red	-16	0.55	0.14
dark	far-red	-12	0.59	0.18

a. Initial curvatures of the hooks average 15 degrees. Final curvatures measured 31 hours after they were excised from the seedling.

The effect of red and far-red light on ethylene evolution and germination of lettuce seeds is shown in Table 2. In this experiment seeds were allowed to imbibe water for 16 hours in the presence of the far-red light source. The bottles were placed 10 to 16 cm from the incandescent light bulb and a fan was placed nearby to reduce the effects of heating. After that time, the bottles were placed either in the dark or given the treatments indicated. Germination and accumulation of ethylene in the gas phase was measured 24 hours later. As shown in Table 2, 1 hour of red light after the 16-hour imbibition period in far-red light stimulated ethylene production. This stimulation was blocked by a terminal far-red exposure and enhanced by a terminal red exposure. Table 2 also shows that ethylene stimulated the germination of seeds receiving both red and far-red light treatments.

TABLE 2. EFFECT OF LIGHT ON ETHYLENE EVOLUTION AND GERMINATION OF GRAND RAPIDS LETTUCE SEEDS

Treatment ^a /	Germination, %	Ethylene in Gas Phase, ^b / ppm
FR	50	1.1
FR-R	82	1.7
FR-R-FR	44	0.8
FR-R-FR-R	86	2.2
FR-R + 10 ppm ethylene	89	-
FR + 10 ppm ethylene	71	-

- a. FR = dark; FR-R = 1-hour exposure to red light; FR-R-FR = additional 1-hour exposure to far-red light; FR-R-FR-R = additional 1-hour exposure to red light.
- b. Bottles were sealed after the last light treatment, and ethylene was measured 24 hours later.

IV. DISCUSSION

Kang et al.⁷ reported that red light inhibited the production of ethylene from bean hypocotyls and that ethylene stimulated hook closing. Our observations confirm this report and suggest that this is a true phytochrome effect because it was reversed by far-red light. Earlier, Goeschl and Pratt¹⁴ reported that morphological responses to light, including the opening of the pea (*Pisum sativum* L. var. Alaska) hook, could be reversed by ethylene. It appears that in hook tissue, ethylene production is inhibited when phytochrome is in the far-red absorbing form and stimulated when it is in the red absorbing form.

The observation that ethylene promotes the germination of seeds has been reported earlier by a number of workers.¹⁵⁻¹⁷ The observation reported here extends this effect of ethylene to another species. As shown in Table 2, a terminal germination-promoting red exposure stimulated ethylene production; a germination-inhibiting far-red exposure inhibited ethylene production. These results suggest that light may regulate ethylene evolution which, in turn, regulates germination.

The method through which phytochrome controls ethylene production is unknown. However, the observation that phytochrome does control ethylene production in bean seedlings and in lettuce seeds suggests possible explanations for other physiological processes. Red light stimulates the formation of pigments associated with ripening in tomato¹⁸ and apple¹⁹ skins. That ethylene does accelerate ripening has been well documented.¹ Red light during the long dark induction period inhibits the flowering of cocklebur (Xanthium pensylvanicum Walln.).²⁰ Recently, ethylene has been shown to have the same effect.³

The number of phytochrome-mediated processes that are mimicked by ethylene remains to be established, but it appears now that this simple hydrocarbon gas acts as an intermediate in some phytochrome-mediated processes as well as in some processes mediated by auxin, herbicide, gibberellin, and abscisin II.

LITERATURE CITED

1. Burg, S.P. 1962. The physiology of ethylene formation. Ann. Rev. Plant Physiol. 13:365-302.
2. Abeles, F.B. 1967. Mechanism of action of abscission accelerators. Physiol. Plant. 20:442-454.
3. Abeles, F.B. 1957. Inhibition of flowering in Xanthium pensylvanicum Walln. by ethylene. Plant Physiol. 42:608-609.
4. Abeles, F.B.; Rubinstein, B. 1964. Regulation of ethylene evolution and leaf abscission by auxin. Plant Physiol. 39:963-969.
5. Burg, S.P.; Burg, E.A. 1966. The interaction between auxin and ethylene and its role in plant growth control. Proc. Nat. Acad. Sci. 55:262-269.
6. Burg, S.P.; Burg, E.A. 1966. Auxin-induced ethylene formation; its relation to flowering in pineapple. Science 152:1268.
7. Kang, B.G.; Yocum, C.S.; Burg, S.P.; Ray, P.M. 1967. Ethylene and carbon dioxide: Mediation of the hypocotyl hook opening response. Science 156:958-959.
8. Zimmerman, P.W.; Wilcoxon, F. 1935. Several chemical growth substances which cause initiation of roots and other responses in plants. Contrib. Boyce Thompson Inst. 7:209-229.
9. Hansen, E. 1946. Effect of 2,4-D on the ripening of Bartlett pears. Plant Physiol. 21:588-592.
10. Maxie, E.C.; Crane, J.C. 1967. 2,4,5-Trichlorophenoxyacetic acid: Effect on ethylene production by fruits and leaves of fig tree. Science 155:1548-1550.
11. Morgan, P.W.; Hall, W.C. 1962. Effect of 2,4-dichlorophenoxyacetic acid on the production of ethylene by cotton and grain sorghum. Physiol. Plant. 15:420-427.
12. Rubinstein, B.; Abeles, F.B. 1965. Relationship between ethylene evolution and leaf abscission. Bot. Gaz. 126:255-259.
13. Klein, W.H.; Withrow, R.H.; Elscud, V.B. 1956. Response of the hypocotyl hook of bean seedlings to radiant energy and other factors. Plant Physiol. 31:289-294.

14. Goeschl, J.D.; Pratt, H.K. 1965. Inhibition by ethylene of some photomorphogenic changes in pea epicotyls. *Plant Physiol.* 40 (Suppl.):111.
15. Vacha, G.A.; Harvey, R.B. 1927. The use of ethylene, propylene, and similar compounds in breaking the rest period of tubers, bulbs, cuttings, and seeds. *Plant Physiol.* 2:187-194.
16. Ruge, V. 1947. Untersuchungen uber Keimungsfordernde Wirkstoffe. *Planta* 35:297-318.
17. Toole, V.K.; Bailer, W.K.; Toole, E.H. 1964. Factors influencing dormancy of peanut seeds. *Plant Physiol.* 39:822-832.
18. Piringer, A.A.; Heinze, P.H. 1954. Effect of light on the formation of a pigment in the tomato fruit cuticle. *Plant Physiol.* 29:467-472.
19. Siegelman, H.W.; Hendricks, S.B. 1958. Photocontrol of anthocyanin synthesis in apple skin. *Plant Physiol.* 33:185-190.
20. Downs, R.J. 1956. Photoreversibility in flower initiation. *Plant Physiol.* 31:279-284.

Unclassified
Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of this, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) Department of the Army Fort Detrick, Frederick, Maryland, 21701		2a. REPORT SECURITY CLASSIFICATION Unclassified	
3. REPORT TITLE PHOTOCONTROL OF ETHYLENE PRODUCTION BY LETTUCE SEEDS AND BEAN HYPOCOTYLS		2b. GROUP	
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)			
5. AUTHOR(S) (First name, middle initial, last name) Abeles, Frederick B. Holm, Robert E. Gahagan, Harry E., III			
6. REPORT DATE July 1967		7a. TOTAL NO. OF PAGES 16	7b. NO. OF REFS 20
8a. CONTRACT OR GRANT NO.		8b. ORIGINATOR'S REPORT NUMBER(S) Technical Manuscript 402	
9. PROJECT NO. IL013001A91A		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
10. DISTRIBUTION STATEMENT Qualified requesters may obtain copies of this publication from DDC. Foreign announcement and dissemination of this publication by DDC is not authorized. Release or announcement to the public is not authorized.			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY Department of the Army Fort Detrick, Frederick, Maryland, 21701	
13. ABSTRACT <p>Red light of about 660 mμ has been shown to promote the germination of <u>Lactuca sativa</u> var. Grand Rapids seeds and to cause the unfolding of the hypocotyl hook of <u>Phaseolus vulgaris</u> L. var. Black Valentine. In both cases, the effect of red light was reversed by far-red light at about 730 mμ.</p> <p>Ethylene promoted the germination of lettuce seeds and caused a negative curvature of bean hooks. Red light stimulated ethylene production from lettuce seeds but it inhibited ethylene production from the bean hooks. Lettuce seed germination and bean hook unfolding seem to be examples in which phytochrome promotes or inhibits the formation of ethylene that, in turn, appears to control the physiological response.</p>			
14. Key Words *Infarfed *Ethylene *Germination Beans Physiology			

DD FORM 1473

REPLACES DD FORM 1473, 1 JAN 64, WHICH IS OBSOLETE FOR ARMY USE.

Unclassified

Security Classification